

To define whether patient is in:
 Acute hyperkalemia[^]
 Chronic hyperkalemia*
 Pseudo-hyperkalemia⁺

eGFR and bicarbonate should also be closely monitored in all hyperkalemia situations


? Pseudo-hyperkalemia

Management

Acute hyperkalemia

Chronic hyperkalemia

Refer to the [KDIGO acute K⁺ management algorithm recommendations](#)



- Pseudo-hyperkalemia is typically defined as a difference of >0.3-0.4 mmol/L between serum and plasma K⁺
- Serum K⁺ measurement should be immediately repeated.
 - Ensure blood is sampled appropriately/or eventually taken as arterial sample.
- In case of hemolysis, to consider whether this occurred in the sample or in the body

^Acute hyperkalemia is defined where a potassium concentration above the upper limit of normal, is not known to be explained by a chronic cause.

***Chronic hyperkalemia** is defined where a potassium concentration above the upper limit of normal, is likely to be explained by a chronic cause (e.g. chronic kidney disease, heart failure, induced by regular medication/supplements), and K⁺ > 5.0mmol/L from repetitive measurements over a 3 month period.

+Pseudo-hyperkalemia is defined where there is a falsely elevated serum potassium concentration, which can occur due to mechanical trauma, prolonged tourniquet use (>1 minute) or fist clenching during the process of blood drawing, and through blood clotting, centrifugation, elevated white blood cell count, or thrombocytosis.



Chronic hyperkalemia Management

Mild (5.0-5.5)

Moderate (>5.5-6.0)

Severe (>6.0)

Important measures to manage hyperkalemia

- Review K⁺ inducing medications and eliminate K⁺ supplements
- See information on [dietary approaches to hyperkalemia in this tool](#)

- Review K⁺ inducing medications and eliminate K⁺ supplements
- See more information on [dietary approaches to hyperkalemia in this tool](#)
- Consider loop diuretics if not prescribed for patients with volume overload, increase loop diuretic dose if already previously prescribed
- Correct acidosis if present

RAASi-specific management

- If on RAASi, aim to maintain RAASi dose and monitor K⁺ levels
- Do not start RAASi if not already prescribed when serum K⁺ >5.0mmol/L
- If indication is for heart failure, consider switch to ARNI from RAASi if available

- Consider K⁺ binder initiation if available to avoid dose reduction
- If on RAASi, and K⁺ binder not available, reduce RAASi dose and monitor K⁺ levels

- Need to reduce K⁺ to <5.0.
- Withhold RAASi and evaluate eGFR, bicarbonate and K⁺ to determine whether RAASi could be restarted
- Consider K⁺ binder initiation if available to facilitate RAASi reinitiation

Additional management in a specific case-to-case basis

- To consider the prescription of SGLT2 inhibitors for patients with eGFR > 25 mL/min/1.73m²